9/28/21, 4:53 PM District Advisory



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

Directorate of Agriculture, Government of Goa,

Panaji



Agromet Advisory Bulletin

Date: 28-09-2021

Weather Forecast of SOUTH-GOA(Goa) Issued On: 2021-09-28(Valid Till 08:30 IST of the next 5 days)

Parameter	2021-09-29	2021-09-30	2021-10-01	2021-10-02	2021-10-03
Rainfall(mm)	34.0	32.0	15.0	7.0	19.0
Tmax(°C)	30.0	31.0	31.0	32.0	36.0
Tmin(°C)	25.0	25.0	25.0	25.0	25.0
RH-I(%)	91	91	92	92	92
RH-II(%)	75	86	81	72	63
Wind Speed(kmph)	14.0	6.0	4.0	4.0	9.0
Wind Direction(Degree)	237	174	120	48	189
Cloud cover(octa)	8	6	6	4	3

Weather Summary/Alert:

• Light to Moderate rainfall/ thundershowers with short intense spells very likely at most places for the district on 28th & 29th Sept 2021 • Light to Moderate rainfall/ thundershowers very likely at most places for the district on 30th Sept 2021 • Light to Moderate rainfall/ thundershowers very likely at many places for the district on 01st & 02nd Oct 2021 • Winds of speed 40 km/hr very likely on 28th & 29th Sept 2021 • Heavy rainfall very likely at isolated places over the district on 28th & 29th Sept 2021 • No weather warnings for the district on 30th Sept, 01st & 02nd Oct 2021 • Maximum & minimum temperatures are likely to be around 30°C & 23°C respectively Week 1 (24th Sep to 30th Sep) • Rainfall amount is very likely to be large excess over North Goa and South Goa districts • Daily monsoon activity is very likely to be normal to active • Maximum and Minimum temperatures are likely to be normal South Goa districts • Daily monsoon activity is very likely to be large excess over North Goa and South Goa districts • Daily monsoon activity is very likely to be normal to active • Maximum and Minimum temperatures are likely to be normal

General Advisory:

• Postpone the application of fertilizer and pesticides for 2 days

SMS Advisory:

• In the view of heavy rainfall warning, make arrangements to drain out excess water from the paddy fields, and plantation crops. Provide proper shelter to livestock and poultry birds

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Make proper arrangements to drain out excess water from the paddy fields

9/28/21, 4:53 PM District Advisory

Crop(Varieties)	Crop Specific Advisory
CASHEW	• Due to forecast of heavy rainfall at isolated places, postpone the spraying of chemicals for 2 days • On cashew build-up of the Tea mosquito bug commences during October with the emergence of new flushes. Hence farmers are advised to look upon the cashew orchards for the Tea mosquito bug incidence • Remove the grubs from the holes with the help of 15 mm chisel and apply Chloropyriphos 20 EC @ 5 ml/liter of water on the stem (swabbing) and drench the same solution @ 2 litre per tree. Apply Neem oil 5 ml/litre of water or neem cake 500g to1 kg / tree

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	• Due to forecast of heavy rainfall at isolated places , postpone the spraying of chemicals for 2 days • The second spilt of fertilizers can be given to Coconut trees of equal to or more than 5 years old. Apply urea 610 g, SSP – 2475 g and Murate of Potash 1200 g per tree • Take up the control measure for bud rot (Spray 1% Bordeaux mixture) • Farmers are suggested to look upon the incidence of Coconut mite and application of Neem cake - $5 kg/palm$ can be done
	• Due to forecast of heavy rainfall at isolated places, postpone the spraying of chemicals for 2 days • The second spilt of fertilizers can be given to Arecanut trees of equal to or more than 5 years old. Apply urea 210 g, SSP – 1100 g and Murate of Potash 200 g per tree • For controlling the of Fruit rot (Koleroga) in Arecanut application of Bordeaux mixture 1% can be done

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	• There is forewarning of very high risk of Babesiosis, for North Goa in the month of October 2021 • Periodical application of acaricides in and around cattle shed is necessary for effective control of ticks • Farmers should approach veterinary hospitals in case of animals showing fever and coffee colored urine • Very high risk of Foot and Mouth Disease (FMD) occurrence in dairy animals • First vaccination at the age of 4 months and thereafter once in six months (Between February – March and August - September) should be followed

Fishieries Specific Advisory:

Fishieries(Varieties)	Fishieries Specific Advisory
CARP	• The water level in the pond should be maintained at least 2 -2.5 m letting the excess water drain out • Removal of unwanted predatory fishes/crabs • Stocking of fish seeds of 80-100 mm size (Catla, Rohu, Common Carp may be followed) at stocking density of 5000 numbers per ha • Feeding of groundnut oil cake: Wheat bran (1:1) ratio at 8-10% of total biomass of the Stock (Total weight of stock) may be followed